



### Jump To:

- [Speech delays and language disorders associated with HIE](#)
- [When should a child get speech-language therapy?](#)
- [What causes speech delays and language disorders?](#)
- [Therapy for speech delays and language disorders](#)
- [What happens during speech language therapy?](#)
- [Additional benefits of speech-language therapy](#)
- [Why is speech-language therapy important?](#)
- [About HIE Help Center](#)

Brain injury due to [hypoxic-ischemic encephalopathy \(HIE\)](#) is rarely confined to a single area of the brain. Because oxygen deprivation affects the connections in the brain on a global level, it is often possible that children with HIE will have multiple interrelated delays in development.



Children with HIE can sometimes have delays in developing speech and language. These delays can sometimes be mitigated, while in other more severe circumstances, children may remain non-verbal and require the use of [alternative or augmentative communication \(AAC\) technologies](#) to potentially assist them in communicating their thoughts, needs, and desires.



Developing a method for communicating helps these children interact with others, develop relationships, learn, work, and socialize.

Speech and language are clearly highly interrelated, but they are not interchangeable (1). Speech refers to the physical act of expressing words and sounds, and encompasses the act of the muscles in the lips, tongue, vocal tract, and jaw that make recognizable sounds. Language, on the other hand, refers to communicating in a systematic and meaningful way.

Because language is related to intelligence, disorders in language acquisition and expression are generally considered more serious than speech disorders. There are two broad categories of language disorders (1):

- Expressive language disorder: The ability to communicate meaning to others
- Receptive language disorder: The ability to understand someone else's message

Language disorders can be expressive, receptive, or mixed receptive-expressive. The signs and symptoms of these disorders can vary depending on the type of disorder a child has.

### Signs of expressive language disorders

Some common signs of expressive language disorders include (2):

- Difficulty constructing grammatically correct or complex sentences
- Difficulty using standard grammar conventions
- Difficulty finding words
- Overuse of placeholder words like 'um' or 'uh'
- Smaller vocabulary than is age-appropriate
- Repetition of certain phrases
- Echoing of questions in responses

### Signs of receptive language disorders



Some common signs of receptive language disorders include (3):

- Difficulty understanding speech
  - Lack of interest or minimal interaction when hearing stories
  - Interrupting others in conversation
  - Difficulty following spoken direction
- Difficulty answering questions, or answers different questions than the ones asked
- Difficulty organizing thoughts

An HIE diagnosis can be difficult, but our team is here to help. Call us when you're ready.

Speech delays and language disorders associated with HIE

There are several conditions that are common in certain children with HIE. Speech and language conditions can be improved via intensive speech-language pathology treatment, and can include:

- **Aphasia:** Aphasia is a language disorder that results from brain damage in a specific part or parts of the brain (4). It can manifest in difficulty writing, reading, understanding, or talking.
- **Dyspraxia:** Dyspraxia is a developmental coordination disorder that affects fine and/or gross motor coordination (5). Some individuals with dyspraxia experience difficulties with coordinating the muscle movements required for spoken language.
- **Dysprosody:** Dysprosody is a disorder that does not affect the language or the words themselves, but the timing, cadence, rhythm, or intensity of the words (6).
- **Dysarthria:** Dysarthria is an abnormal facial muscle tone condition that results from brain damage (7). It affects muscles in the mouth, lips, tongue, or throat used for speech.
- **Fluency Disorders:** A speech disorder that relates to rate, smoothness, continuity, or effort in speech (8). The most common fluency disorder is stuttering.
- **Dysphagia:** Dysphagia is a condition that involves difficulty swallowing (9). It can manifest



in many different ways, but involves general pain and hoarseness when speaking.

- Resonance Disorders: Resonance disorders involve abnormalities in the vocal tract that affect the sound of speech (10).

### When should a child get speech-language therapy?

If a child is not acquiring speech or language at the usual rate, it is recommended that parents work with the child's pediatrician to have the child evaluated *as soon as possible* (during infancy or very early childhood). Sometimes, children may require some extra assistance or therapy to help them acquire language, or may need extra intervention in the case of concurrent developmental issues.

It is critical to make these diagnoses early, so the child may be enrolled in an Early Intervention (Zero to Three, Head Start, Early On) program to mitigate some of the difficulties that may be associated with poor communication. These can include future academic struggles, emotional or behavioral problems associated with poor social interaction, and difficulties in occupational or psychological development. These programs are most effective when a child is very young, as beginning these programs at an older age will mean that the child must also take the extra step of unlearning old frameworks and developing new ones.

It is also worth noting that sometimes speech and/or language delays can result from hearing-related deficits. Hearing problems may make it appear that a child has a language or speech issue when they only require a hearing aid or medical intervention to get their speech and/or language development on track. To learn more about hearing issues associated with HIE, click [here](#).

### What causes speech delays and language disorders?

Delays in speech and/or language can be due to several root causes. In some children, HIE-related speech/language disorders are due to damage to the part of the brain responsible for processing language, which can require a different approach from speech/language disorders arising from damage to parts of the brain that control the muscles of the throat, mouth, and



jaw (a common feature of [cerebral palsy](#) stemming from HIE).

A licensed speech-language therapist will be able to determine the best way to assist a child with a speech/language disorder, working with the child's caregiver team of physicians, physical therapists, psychologists, and social workers to help the child overcome cognitive or developmental difficulties that can make communication difficult.

### Therapy for speech delays and language disorders

Usually, when a child is evaluated and found to have a speech or language disorder, they can be referred to a speech and language pathologist. A speech-language pathologist is an American Speech-Language-Hearing Association (ASHA)-certified professional (with a minimum of a Master's degree in speech and language pathology) that helps individuals with difficulty communicating adapt and compensate to maximize their communication abilities. These professionals have training in pathology, audiology, anatomy and physiology, linguistics and phonetics, human development, and American Sign Language (ASL), among other courses, which allows them to effectively assist children with specialized developmental needs.

Parents seeking a speech-language pathologist for their children can request assistance at a variety of locations, as speech-language pathologists are often available at clinics, community health centers, hospitals, rehabilitation centers, public schools, nursing homes and state health departments.

In some circumstances, children receive one-on-one services with additional exercises that they practice at home with parents outside of therapy sessions. In other instances, speech-language pathology services are provided in a small group setting. Sometimes, these services can be provided in a home setting without having to travel.

What happens during speech language therapy?



First and foremost, it is important to remember that each child is different, so the exercises that a speech-language pathologist recommends may differ substantially from child to child. For children who are verbal, speech-language therapy can help them (11):

- Pronounce words more clearly
- Listen to and understand conversation
- Use words properly
- Understand grammar
- Have a bigger vocabulary
- Comprehend conversation
- Speak in complex sentences
- Speak conversationally
- Exercise memory and recall
- Understand the differences and similarities between written text and spoken words

A speech-language pathologist will not only help the child physically form words needed for communication but will also develop a training regimen designed to help the child's brain interpret and understand words, sounds, numbers and gestures.

Some children with HIE may be non-verbal. When working with children who are nonverbal, speech-language pathologists can do the following (12):

- Create and encourage alternative communication strategies, such as gestures, sign language, writing, and others
- Improve social skills through role playing, social stories, and other tools
- Focus teaching on the language, through comprehension, sequencing, and other tools
- Aid in communication through reading

Additional benefits of speech-language therapy

Speech-language therapy revolves around the use of the muscles involved in the functioning of the throat, mouth, and jaw. While communication is the primary goal of speech-language



therapy, it often has the added benefit of helping children drink, eat, breathe and chew more effectively as well. The therapy can be catered to a child's specific needs. Often, children with HIE who have cerebral palsy will have a degree of impairment in eating and drinking (dysphagia); speech-language pathologists can assist these children in improving complex muscle coordination. This decreases the risk that a child will aspirate their food (which can lead to choking and/or pneumonia), have abnormal breathing, increased coughing, dehydration, or malnutrition.

### Why is speech-language therapy important?

Communication is the underpinning of social interaction, which makes early intervention critical. Developing a communication system helps to ensure that the child will have frameworks in place for inclusion and socialization, allowing them to more fully engage with their surroundings, participate in activities in school or at home, learn new skills, and develop new hobbies or interests.

This therapy also increases the likelihood that other interventions will be effective. Because other types of therapies are dependent on the child's ability to comprehend and follow instruction, speech-language therapy is a crucial part of a child's multidisciplinary therapy regimen. Speech-language therapy supports learning, education, literacy, and community integration.

There are no risks to speech-language therapy. Delaying speech-language pathology can increase the risk of isolation or depression, and can mean that other areas of development (social, educational, psychological, etc.) may also slow without continued communication-related support.

### Additional speech and language-related resources

- [University of Michigan: Speech and Language Delay and Disorder Quick Reference](#)
- [What Does Normal Speech Acquisition Look Like?](#)



- [Speech Therapy: How it Works, What to Expect](#)
- [How Long Does it Take For Speech Therapy to Work?](#)
- [Encyclopedia of Children's Health: Language Delays](#)
- [A Parent's Guide to Speech and Language Disorders](#)

### About the HIE Help Center and ABC Law Centers

The HIE Help Center is run by [ABC Law Centers](#), a medical malpractice firm exclusively handling cases involving HIE and other birth injuries. Our lawyers have over 100 years of combined experience with this type of law, and have been advocating for children with HIE and related disabilities since the firm's inception in 1997.

We are passionate about helping families obtain the compensation necessary to cover their extensive medical bills, loss of wages (if one or both parents have to miss work in order to care for their child), assistive technology, and other necessities.

If you suspect your child's HIE may have been caused by medical negligence, please [contact us](#) today to learn more about pursuing a case. We provide free legal consultations, during which we will inform you of your [legal options](#) and answer any questions you have. Moreover, you would pay nothing throughout the entire legal process unless we obtain a favorable settlement.

You are also welcome to reach out to us with inquiries that are not related to malpractice. We cannot provide individualized medical advice, but we're happy to track down informational resources for you.

### Sources

Koslo, E. (n.d.). What's the Difference Between a Speech Impairment and a Language Disorder? Retrieved June 28, 2019, from <https://www.understood.org/en/learning-attention-issues/child-learning-disabilities/communication-disorders/difference-between-speech-impairment-and-language-disorder>





Developmental Expressive Language Disorder (DELD). (n.d.). Retrieved June 28, 2019, from <https://www.healthline.com/health/expressive-language-disorder-developmental#symptoms>

Rosen, P. (n.d.). Receptive Language Issues: What You're Seeing. Retrieved June 28, 2019, from <https://www.understood.org/en/learning-attention-issues/child-learning-disabilities/communication-disorders/receptive-language-issues-what-youre-seeing>

Aphasia. (n.d.). Retrieved June 28, 2019, from <https://www.asha.org/public/speech/disorders/Aphasia/>

Speech and Language. (2013, October 07). Retrieved June 28, 2019, from <https://dyspraxiafoundation.org.uk/about-dyspraxia/speech-language/>

(n.d.). Retrieved June 28, 2019, from <http://www.speechdisorder.co.uk/dysprosody.html>

Dysarthria. (n.d.). Retrieved June 28, 2019, from <https://www.asha.org/public/speech/disorders/dysarthria/>

Childhood Fluency Disorders. (n.d.). Retrieved June 28, 2019, from <https://www.asha.org/practice-portal/clinical-topics/childhood-fluency-disorders/>

Dysphagia. (2018, February 03). Retrieved June 28, 2019, from <https://www.mayoclinic.org/diseases-conditions/dysphagia/symptoms-causes/syc-20372028>

Resonance Disorders. (n.d.). Retrieved June 28, 2019, from <https://www.asha.org/Practice-Portal/Clinical-Topics/Resonance-Disorders/>

Who Are Speech-Language Pathologists, and What Do They Do? (n.d.). Retrieved July 1, 2019, from <https://www.asha.org/public/Who-Are-Speech-Language-Pathologists/>

5 Reasons Why Speech & Language Therapy is Crucial for a Non-Verbal Child – Friendship Circle – Special Needs Blog. (2015, May 14). Retrieved June 28, 2019, from <https://www.friendshipcircle.org/blog/2015/05/14/5-reasons-why-speech-language-therapy-is-crucial-for-a-non-verbal-child/>